

# Western Australian Immunisation Schedule

## Childhood

Age and eligibility	Antigen/Antibody	Brand(s)	Important information
<b>Birth to &lt;7 days</b>	Nirsevimab*	Beyfortus®	Can be given up to 8 months of age, but preferably as close to birth as possible, if the mother did not receive Abrysvo > 2 weeks prior to birth. See RSV table and links below for further information.
	Hepatitis B	H-B-Vax® II Paed or Engerix® B Paed	
<b>6-8 weeks</b>	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
	13vPCV	Prevenar® 13	
	Rotavirus	Rotarix®	Oral dose 1: Must be given by 14 weeks of age
<b>Aboriginal*and/or MRC</b>	MenACWY	Nimenrix®	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero®	<a href="#">Prophylactic paracetamol</a> recommended
<b>4 months</b>	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
	13vPCV	Prevenar® 13	
	Rotavirus	Rotarix®	Oral dose 2: Must be given by 24 weeks of age
<b>Aboriginal*and/or MRC</b>	MenACWY	Nimenrix®	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero®	<a href="#">Prophylactic paracetamol</a> recommended
<b>6 months</b>	DTPa-hepB-IPV-Hib	Infanrix® hexa or Vaxelis®	
<b>Aboriginal and/or MRC</b>	13vPCV	Prevenar® 13	
<b>MRC</b>	MenB	Bexsero®	<a href="#">Prophylactic paracetamol</a> recommended
	MenACWY	Nimenrix®	
<b>12 months</b>	MMR	Priorix® or M-M-R II®	The 2 brands of MMR vaccine are interchangeable
	13vPCV	Prevenar® 13	
	MenACWY	Nimenrix®	
<b>Aboriginal and/or MRC</b>	MenB	Bexsero®	<a href="#">Prophylactic paracetamol</a> recommended
<b>MRC</b>	Hepatitis B	H-B-Vax® II Paed or Engerix® B Paed	<32 weeks gestation or <2000g birth weight
<b>18 months</b>	DTPa	Infanrix® or Tripacel®	
	Hib	ActHIB®	
	MMRV	Priorix-Tetra®	Do not administer MMRV as dose 1 of a measles-containing vaccine if <4 years of age
<b>Aboriginal</b>	Hepatitis A	Vaqa® Paed/Adol	Dose 1 of 2. Dose 2 to be administered at least 6 months after dose 1
<b>&lt;2 years</b>	Catch-up for MenB is available for Aboriginal children <2 years of age		
<b>Aboriginal</b>			
<b>4 years</b>	DTPa-IPV	QuadraceI® or Infanrix® IPV	
<b>Aboriginal</b>	Hepatitis A	Vaqa® Paed/Adol	Not required if previously received 2 doses (first dose at age ≥12 months) at least 6 months apart
<b>Aboriginal and/or MRC</b>	23vPPV	Pneumovax23®	Dose 1 (if not previously administered). Dose 2 should be administered at least 5 years later

## Respiratory Syncytial Virus (RSV)

Age and eligibility	Antigen/Antibody	Brand(s)	Important information
<p>See here for WA state-wide eligibility, excluding Kimberley/ Pilbara regions.</p> <p>See here for Kimberley and Pilbara regions eligibility only.</p>	Nirsevimab*	Beyfortus®	The RSV Immunisation decision aid contains information on both Abrysvo for patients currently pregnant, and Beyfortus for children under 2 years of age.

## Adolescent and adult

Age and eligibility	Antigen	Vaccine brand(s)	Important information
Year 7	HPV	Gardasil®9	Catch-up for adolescents <26 years of age
	dTpa	Boostrix® or Adacel®	
Year 10	MenACWY	Nimenrix® or MenQuadfi®	
≥18 years MRC	Herpes-zoster	Shingrix®	2 doses, given 1–2 months apart
Pregnant	dTpa	Boostrix® or Adacel®	Each pregnancy, ideally between 20–32 weeks
	RSV	Abrysvo®	Eligible women between 28 - 36 weeks
≥20 years Aboriginal*	Hepatitis B	Engerix® B	For non-immune persons. See the <a href="#">Australian Immunisation Handbook</a> for recommendations and timing of doses
Adult refugees and humanitarian entrants	dTpa (Boostrix® or Adacel®), HepB (H-B-Vax® II or Engerix® B), IPV (IPOL®), MMR (MMRII® or Priorix®), VZV (Varivax®). HPV (Gardasil®9) <26 years of age		See the <a href="#">Australian Immunisation Handbook</a> for timing of doses
≥50 years Aboriginal	13vPCV 23vPPV	Prevenar® 13 Pneumovax23®	See the <a href="#">Australian Immunisation Handbook</a> for timing of doses
	Herpes-zoster	Shingrix®	2 doses, given 2–6 months apart
Born after 1965*	MMR	Priorix® or M-M-R II®	Give 1 or 2 doses as required. Whilst serology is not routinely recommended, see <a href="#">WA adult measles vaccination program</a> for details. The 2 brands of MMR vaccine are interchangeable
≥65 years	Herpes-zoster	Shingrix®	2 doses, given 2–6 months apart
≥70 years	13vPCV	Prevenar® 13	

All individuals diagnosed with **specified MRC** aged >12 months of age require 1 dose of 13vPCV at diagnosis, followed by a first dose of 23vPPV, 2 to 12 months later or at 4 years of age, whichever is later. A second dose is given at least 5 years later

Additional *Haemophilus influenzae* type b (Hib), meningococcal and pneumococcal vaccines are recommended for people with specified risk conditions. Refer to the [Australian Immunisation Handbook](#) for details

Age and eligibility	Vaccine brand(s)	Important information
6 Months to <5 Years	Vaxigrip®Tetra, FluQuadri®	For children <9 years, or persons following a stem cell or solid organ transplant, 2 doses are required 1 month apart in the first year they receive influenza vaccine. One dose annually in subsequent years. For information on influenza dosing see the <a href="#">Australian Immunisation Handbook</a> or annual <a href="#">ATAGI advice</a> on seasonal influenza vaccines.
≥5 Years to Year 6*	FluQuadri®	
5 Years to <65 Years	Flucelvax®Quad, Vaxigrip®Tetra	
Aboriginal and/or MRC	Flucelvax®Quad, Vaxigrip®Tetra	
Vulnerable cohorts* 12 to <65 Years	FluQuadri®	For non-NIP eligible residents in congregate living and people experiencing homelessness.
Pregnancy	Vaxigrip®Tetra, Flucelvax®Quad	Each pregnancy, any trimester
All West Australians 12 to <65 years*	FluQuadri®	Eligibility under Free Influenza Program May, June and July 2025
≥65 Years	Fluad® Quad	Preferentially use adjuvanted QIV
WA Health staff^	Flucelvax®Quad, FluQuadri®, Fluad® Quad	For ≥65 years use Fluad® Quad

\*Refers to WA State-funded Programs.

^Refers to all persons attending WA Health staff influenza vaccination services.

The term '**Aboriginal**' is inclusive of all Aboriginal and Torres Strait Islander people.

**MRC** refers to people with medical risk conditions. Refer to the current online edition of the [Australian Immunisation Handbook](#) for all medical risk conditions, recommendations and timing of doses.

**Persons aged <20 years** can catch-up scheduled vaccines, refer to [Australian Immunisation Handbook](#) for timing of doses.